



# WAKULLA SPRINGS AND RIVER

## 2016 GRADE


## C




## A




## D




## C



## D




## F




## B

# WAKULLA SPRINGS AND RIVER





<b>Spring Discharge</b>	<b>Grade</b>	<b>Average (2012-2016)</b>	<b>A: &gt; 654 cfs</b> <b>B: 493 - 653 cfs</b> <b>C: 364 - 492 cfs</b> <b>D: 290 - 363 cfs</b> <b>F: &lt; 289 cfs</b>
	<b>A</b>	<b>805 cfs</b>	

# WAKULLA SPRINGS AND RIVER




<b>Water Clarity @ Spring Run</b>	<b>Grade</b>	<b>Average (2015-2016)</b>	<b>A: &gt; 70 %</b>
	<b>D</b>	<b>49%</b> Light Transmittance @ 1 meter	<b>B: 60 - 69 %</b> <b>C: 50 - 59 %</b> <b>D: 40 - 49 %</b> <b>F: &lt; 40 %</b>

# WAKULLA SPRINGS AND RIVER




Nitrate Concentration @ Spring Boil	Grade	Average (2016)	A: < 0.15 mg/L B: 0.15 - 0.35 mg/L C: 0.36 - 0.60 mg/L D: 0.61 - 0.90 mg/L F: > 0.90 mg/L
	<b>C</b>	<b>0.45 mg/L</b>	

# WAKULLA SPRINGS AND RIVER




<b>Adult Aquatic Emergence @ Spring Run</b>	<b>Grade</b>	<b>Average (2015-2016)</b>	<b>A: &gt; 100</b> <b>B: 75 - 99</b> <b>C: 50 - 74</b> <b>D: 25 - 49</b> <b>F: &lt; 24</b>
	<b>D</b>	<b>30 insects/m<sup>2</sup>/day</b>	

# WAKULLA SPRINGS AND RIVER



Fish Biomass	Grade	Average (2016)	<b>A: &gt; 300</b> <b>B: 200 - 299</b> <b>C: 100 - 199</b> <b>D: 50 - 99</b> <b>F: &lt; 50</b>
	<b>F</b>	<b>5.9</b> Biomass kg/ha	

# WAKULLA SPRINGS AND RIVER



<b>Photosynthetic Efficiency</b>	<b>Grade</b>	<b>Average (2015-2016)</b>	<b>A: &gt; 7.0</b> <b>B: 5.0 - 6.99</b> <b>C: 3.0 - 4.99</b> <b>D: 1.0 - 2.99</b> <b>F: &lt; 1.0</b>
	<b>B</b>	<b>5.92 %</b>	